

Ibuprofen Dosing Chart

(Also known as Motrin®, Advil®)

Age	Weight	Ibuprofen Infant Drops	Ibuprofen Suspension	Ibuprofen Children's Chewable	Ibuprofen Children's Chewable
		50 mg/1.25 ml	100 mg/5 ml	50 mg ea.	100 mg ea.
6-11 mo.	12-17 lbs.	50 mg/.1.25 ml			
12-23 mo.	18-23 lbs.	75 mg/1.8 ml			
2-3 yrs.	24-35 lbs.		100 mg/1 tsp.	100 mg/2 tabs	100 mg/ 1 tab
4-5 yrs.	36-47 lbs.		150 mg/1.5 tsp.	150 mg/3 tabs	150 mg/1.5 tabs
6-8 yrs.	48-59 lbs.		200 mg/2 tsp.	200 mg/4 tabs	200mg/2 tabs
9-10 yrs.	60-71 lbs.		250 mg/2.5 tsp.	250 mg/5 tabs	250 mg/2.5 tabs
11 yrs.	72-95 lbs.		300 mg/3 tsp.	300mg/6 tabs	300 mg/3 tabs

Do not give more frequently than every 4 hours or more than 5 doses in 24 hours.

Never alternate dosing of Ibuprophen and Acetaminophen unless your doctor tells you to. Start with one medication and stay with it. Ibuprofen (Children's Advil or Motrin) can be given to children six months and older. However, do not give to a child who has been drinking poorly, is nauseous, vomiting, has stomach pains, a history of stomach ulcer, chicken pox, is undergoing chemotherapy or has bleeding

When Should I Call A Doctor?

- When an infant under three (3) months has a fever, is irritable or cries persistently.
- If a child complains of a stiff neck, headache or severe pain or has a seizure.
- If a child shows signs of having a difficult time breathing.
- When a child develops a persistent fever.
- When you have concerns or questions.